



## MEDICATION ADMINISTRATION AUTHORIZATION FORM

2024/2025

Medication must be delivered to school by a responsible adult (whenever possible, a parent) in the original pharmacy labeled container or unopened manufacturer's package. Medication may not be delivered directly by the student. Medication is counted and documented upon receipt and signed by the school staff member and adult delivering the medication. This form must be renewed annually and must be signed by a non-parent, licensed prescriber.

### LICENSED PRESCRIBER'S AUTHORIZATION

Name of Student: \_\_\_\_\_ Date of Birth: \_\_\_\_\_ Grade: \_\_\_\_\_

Condition for which medication is being administered: \_\_\_\_\_

Medication Name: \_\_\_\_\_

Dose: \_\_\_\_\_ Route: \_\_\_\_\_ Time(s): \_\_\_\_\_

If PRN, frequency: \_\_\_\_\_ If PRN, for what symptoms: \_\_\_\_\_

Relevant side effects: ☐ None expected ☐ Specify: \_\_\_\_\_

Medication shall be administered from: \_\_\_\_\_ to \_\_\_\_\_

Month / Day / Year

Month / Day / Year

Prescriber's Name/Title \_\_\_\_\_

Telephone: \_\_\_\_\_ FAX: \_\_\_\_\_

Address: \_\_\_\_\_

Licensed Prescriber's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

(Original signature or signature stamp ONLY)

### PARENT/GUARDIAN AUTHORIZATION

I/We request designated school personnel to administer the medication as prescribed by the above prescriber. I/We certify that I/we have legal authority to consent to medical treatment for the student named above, including the administration of medication at school. I/We understand that within one week of the end of the school year, an adult must pick up the medication, otherwise it will be discarded. I/We authorize the school administration to communicate with the health care provider as allowed by HIPAA.

Parent/Guardian Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Home #: \_\_\_\_\_ Cell #: \_\_\_\_\_ Work #: \_\_\_\_\_

### EMERGENCY MEDICINE SELF CARRY/SELF ADMINISTRATION AUTHORIZATION

Student may self-carry this emergency medication. (must be signed by prescriber and principal)

Licensed Prescriber's Signature \_\_\_\_\_

Principal's Signature \_\_\_\_\_ Date \_\_\_\_\_

I understand point 12e of the Medication Administration Parent Procedure (Form MED-1) and agree to abide by it.

Student's Signature \_\_\_\_\_ Date \_\_\_\_\_

Medication Administration Monitor Approval: \_\_\_\_\_

GCS Form  
MED-4 (2/24)



## EMERGENCY MEDICATION SELF-CARRY AND SELF-ADMINISTRATION CARE PLAN

2024-2025

I, \_\_\_\_\_, agree to keep record of when I self-administer my medication. I understand that I may need to share this record with my parents or the Medication Administration Monitor if requested.

Medication name \_\_\_\_\_

Dosage \_\_\_\_\_

Time to be taken \_\_\_\_\_

In the event that I am unable to self-administer my medication, students and staff around me should \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_

911 should be called if \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_

Student signature \_\_\_\_\_ Date \_\_\_\_\_

Parent signature \_\_\_\_\_ Date \_\_\_\_\_

"MAMi" signature \_\_\_\_\_ Date \_\_\_\_\_

Principal signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

Teacher signature \_\_\_\_\_ Date \_\_\_\_\_

[illegible]

Name: \_\_\_\_\_ D.O.B.: \_\_\_\_\_

Allergic to: \_\_\_\_\_

 Weight: \_\_\_\_\_ lbs. Asthma: ☐ **Yes (higher risk for a severe reaction)** ☐ **No**

**PLACE  
PICTURE  
HERE**

**NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.**

**Extremely reactive to the following allergens:** \_\_\_\_\_

**THEREFORE:**

- ☐ If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- ☐ If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR **ANY** OF THE FOLLOWING:  
**SEVERE SYMPTOMS**



**LUNG**

Shortness of breath, wheezing, repetitive cough



**HEART**

Pale or bluish skin, faintness, weak pulse, dizziness



**THROAT**

Tight or hoarse throat, trouble breathing or swallowing



**MOUTH**

Significant swelling of the tongue or lips



**SKIN**

Many hives over body, widespread redness



**GUT**

Repetitive vomiting, severe diarrhea



**OTHER**

Feeling something bad is about to happen, anxiety, confusion

**OR A  
COMBINATION**  
of symptoms  
from different  
body areas.

1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
  - Consider giving additional medications following epinephrine:
    - » Antihistamine
    - » Inhaler (bronchodilator) if wheezing
  - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
  - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
  - Alert emergency contacts.
  - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

## MILD SYMPTOMS



**NOSE**

Itchy or runny nose, sneezing



**MOUTH**

Itchy mouth



**SKIN**

A few hives, mild itch



**GUT**

Mild nausea or discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE  
SYSTEM AREA, GIVE EPINEPHRINE.**

**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM  
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

## MEDICATIONS/DOSES

Epinephrine Brand or Generic: \_\_\_\_\_

Epinephrine Dose: ☐ 0.1 mg IM ☐ 0.15 mg IM ☐ 0.3 mg IM


Antihistamine Brand or Generic: \_\_\_\_\_

Antihistamine Dose: \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if wheezing): \_\_\_\_\_


## HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




## HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



## HOW TO USE SYMJEPI™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPI by finger grips only and slowly insert the needle into the thigh. SYMJEPI can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



## ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

**OTHER DIRECTIONS/INFORMATION** (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

## EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: \_\_\_\_\_

DOCTOR: \_\_\_\_\_ PHONE: \_\_\_\_\_

PARENT/GUARDIAN: \_\_\_\_\_ PHONE: \_\_\_\_\_

## OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: \_\_\_\_\_ PHONE: \_\_\_\_\_

NAME/RELATIONSHIP: \_\_\_\_\_ PHONE: \_\_\_\_\_

NAME/RELATIONSHIP: \_\_\_\_\_ PHONE: \_\_\_\_\_